Neuroscience Letters, 92 (1988) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abbenante, J., see Kerr, D.B., 92
Aguayo, L.G. and Weight, F.F., Membrane currents activated in acutely dissociated rat pineal cells during the circadian cycle, 155
Akasu, T., 5-Hydroxytryptamine facilitates GABA-induced depolarization in bullfrog primary afferent neurons, 270
Åkerman, K.E.O., Enkvist, M.O.K. and Holopainen, I., Activators of protein kinase C and phenylephrine depolarize the astrocyte membrane by reducing the K^+ permeability, 265
Alford, S.T., see Davies, S.N., 213
Allsop, D., Ikeda, S., Bruce, M. and Glenner, G.G., Cerebrovascular amyloid in scrapie-affected sheep reacts with antibodies to prion protein, 234

Balayadi, M., Jule, Y. and Cupo, A., Enkephalin-like immunoreactive principal ganglion cells and nerve fibres in inferior mensenteric ganglion of the cat, 126
Baptista, T., Parada, M.A. and Murzi, E., Puberty modifies sulpiride effects on body weight in rats, 161
Barnes, S. and Davies, J.A., The effects of calcium channel agonists and antagonists on the release of endogenous glutamate from cerebellar slices, 58
Beattie, D.T., see Curtis, D.R., 97
Béloeil, J.C., see Jacquin, T., 285
Ben-Ari, Y., see Neuman, R.S., 64
Bhattacharya, S.K., Glover, V., McIntyre, I., Oxenkrug, G. and Sandler, M., Stress causes an increase in endogenous monoamine oxidase inhibitor (tribulin) in rat brain, 218
Bonhoeffer, T. and Staiger, V., Optical recording with single cell resolution from monolayered slice cultures of hippocampus, 259
Briotta, G.D., see Hösli, L., 165
Bruce, M., see Allsop, D., 234
Bruno, J.P., see Weihmuller, F.B., 207
Burnstock, G., see Fehér, E., 137
Bush, K.T., see Nagele, R.G., 241

Carrer, H.F., see Ramírez, O.,A., 275
Ceballos, I., see Delacourte, A., 247
Champagnat, J., see Jacquin, T., 285
Chen, M., see Yinon, U., 280
Cherubini, E., see Neuman, R.S., 64
Cheung, W.-T., see Lee, C.-M., 310
Coan, E.J., see Davies, S.N., 213
Collingridge, G.L., see Davies, S.N., 213
Cordo, P.J., Kinesthetic coordination of a movement sequence in humans, 40
Coutable, D., see Sarrieau, A., 14
Cupo, A., see Balayadi, M., 126
Curtis, D.R., Gynther, B.D., Beattie, D.T., Kerr, D.I.B. and Prager, R.H., Baclofen antagonism by 2-hydroxy-saclofen in the cat spinal cord, 97

Dall'Olio, R., see Gandolfi, O., 192
Daval, J.L., Deckert, J., Nakajima, T., Morgan, P.F. and Marangos, P.J., Regional ontogenetic profile of central and peripheral benzodiazipine receptors in the guinea pig brain, 82
Davies, J.A., see Barnes, S., 58
Davies, S.N., Alford, S.T., Coan, E.J., Lester, R.A.J. and Collingridge, G.L., Ketamine blocks an NMDA receptor-mediated component of synaptic transmission in rat hippocampus in a voltage-dependent manner, 213
De Caro, G., see Massi, M., 341
Deckert, J., see Daval, J.L., 82
Defossez, A., see Delacourte, A., 247
Del Bianco, E., see Santicioli, P., 197
Del Cerro, C.D., see Del Cerro, M., 21
Del Cerro, M., Notter, M.F.D., Wiegand, S.J., Jiang, L.Q. and Del Cerro, C.D., Intraretinal transplantation of fluorescently labeled retinal cell suspensions, 21
Delacourte, A., Defossez, A., Ceballos, I., Nicole, A. and Sinet, P.M., Preferential localization of copper zinc superoxide dismutase in the vulnerable cortical neurons in Alzheimer's disease, 247
Dillon, D.L., see McRae-Degueurce, A., 303
Doerffler-Melly, J. and Neuhuber, W.L., Rectospinal neurons: evidence for a direct projection from the enteric to the central nervous system in the rat, 121
Doherty, P., Seaton, P., Flanigan, T.P. and Walsh, F.S., Factors controlling the expression of the NGF receptor in PC12 cells, 222
Dussaillant, M., see Sarrieau, A., 14

Enkvist, M.O.K., see Åkerman, K.E.O., 265
Erdélyi, L. and Such, G., The A-type potassium current: catechol-induced blockage in snail neurons, 46

Fehér, E. and Burnstock, G., Galanin-like immunoreactive nerve elements in the small intestine of the rat. An electronmicroscopic immunocytochemical study, 137
Flanigan, T.P., see Doherty, P., 222
Fortin, G., see Jacquin, T., 285
Fujii, T., see Kunugi, Y., 315
Fujiki, N., see Hirayama, M., 34

Gandolfi, O., Roncada, P. and Dall'Olio, R., Single or repeated administration of SCH 23390 fail to affect serotonergic neurotransmission, 192
Gavras, H., see Vlahakos, D., 335
Ge, M., Ueda, H. and Satoh, M., Endogenous GABA released into the fourth ventricle of the rat brain in vivo is enhanced by noxious stimuli, 76
Gelerstein, S., see Yinon, U., 280
Gentili, L., see Massi, M., 341
Geppetti, P., see Santicioli, P., 197
Gho, M., see Neuman, R.S., 64
Gillet, B., see Jacquin, T., 285
Glenner, G.G., see Allsop, D., 234
Glover, V., see Bhattacharya, S.K., 218
Gonzalez-Brito, A., Reiter, R.J., Tannenbaum, M.G. and Jones, D.J., Adrenalectomy but not gonadectomy affects rat pineal β-adrenergic receptor density, 330
Greenstein, Y.J., see Pavlides, C., 177
Gynther, B.D., see Curtis, D.R., 97

Haas, H.L., see Regenold, J.T., 347
Hadjiconstantinou, M., see Weihmuller, F.B., 207
Hashimoto, Y., see Kunugi, Y., 315
Heuss, L., see Hösli, L., 165
Hirayama, M., Miyadai, T., Yokochi, T., Sato, K., Kubota, T., Iida, M. and Fujiki, N., Infection of human T-lymphotropic virus type I to astrocytes in vitro with induction of the class II major histocompatibility complex, 34
Hjorth, S., see McRae-Degueurce, A., 303
Holopainen, I., see Åkerman, K.E.O., 265
Homma, I., see Iwase, M., 30
Hornykiewicz, O., see Pifl, Ch., 228
Hösli, L., Briotta, G.D., Quadri, L. and Heuss, L., Action of acetylcholine, muscarine, nicotine and antagonists on the membrane potential of astrocytes in cultured rat brainstem and spinal cord, 165
Hughes, S. and Smith, M.E., Effect of nerve section on β-endorphin and α-melanotropin immunoreactivity in motor nerves of normal and dystrophic mice, 1
Hydman, A.G., see Zeevalk, G.D., 149

Iida, M., see Hirayama, M., 34
Ikeda, S., see Allsop, D., 234
Illes, P., see Regenold, J.T., 347
Inagaki, C., see Kunugi, Y., 315
Iwase, M., Homma, I., Shioda, S. and Nakai, Y., Thyrotropin-releasing hormone-like iummunoreactive neurons in rabbit medulla oblongata, 30

Jackson, D.A. and White, S.R., Thyrotropin releasing hormone (TRH) modifies excitability of spinal cord dorsal horn cells, 171
Jacquin, T., Fortin, G., Pasquier, C., Gillet, B., Béloeil, J.C. and Champagnat, J., Metabolic acidosis induced by *N*-methyl-D-aspartate in brain slices of the neonatal rat: ^{31}P- and ^{1}H-magnetic resonance spectroscopy, 285
Jiang, L.Q., see Del Cerro, M., 21
Johnston, G.A.R., see Kerr, D.B., 92
Jones, D.J., see Gonzalez-Brito, A., 330
Jule, Y., see Balayadi, M., 126

Kerr, D.B., Ong, J., Johnston, G.A.R., Abbenante, J. and Prager, R.H., 2-Hydroxy-saclofen: an improved antagonist at central and peripheral $GABA_B$ receptors, 92
Kerr, D.I.B., see Curtis, D.R., 97
Kijima, H., see Tanabe, N., 52
Klareskog, L., see Maehlen, J., 8
Komuro, T., Direct contacts between Auerbach's ganglia and smooth muscle cells in the small intestine of the rat, 27
Kristensson, K., see Maehlen, J., 8
Kubota, T., see Hirayama, M., 34
Kunugi, Y., Hashimoto, Y., Tamura, A., Fujii, T and Inagaki, C., Effects of phospholipases on Cl-ATPase in the rat brain, 315
Kuraishi, Y., Nanayama, T., Ohno, H., Minami, M. and Satoh, M., Antinociception induced in rats by intrathecal administration of antiserum against calcitonin gene-related peptide, 325

Leanza, G., Pellitteri, R., Russo, A. and Stanzani, S., Branching axons from subcoeruleus area project to the nucleus raphe pontis and hypothalamic zona incerta, as studied with the double fluorescent retrograde tracing technique, 132
Lee, C.-M. and Cheung, W.-T., Effects of neonatal monosodium glutamate treatment on substance

P binding sites in the rat retina, 310
Lee, H.-Y., see Nagele, R.G., 241
Lehman, J., see Tsai, C., 298
Lester, R.A.J., see Davies, S.N., 213
Lothman, E.W., see Stringer, J.L., 69
Lovick, T.A., GABA-mediated inhibition in nucleus paragigantocellularis lateralis in the cat, 182

Maehlen, J., Schröder, H.D., Klareskog, L., Olsson, T. and Kristensson, K., Axotomy induces MHC class I antigen expression on rat nerve cells, 8
Maggi, C.A., see Massi, M., 341
Maggi, C.A., see Renzi, D., 254
Maggi, C.A., see Santicioli, P., 197
Majocha, R.E., see Shea, T.B., 291
Marangos, P.J., see Daval, J.L., 82
Marotta, C.A., see Shea, T.B., 291
Mason, D.W., see McRae-Degueurce, A., 303
Mason, R., Altered neuronal sensitivity of lateral geniculate neurones to noradrenaline and 5-HT following exposure to continuous lighting, 102
Massi, M., Polidori, C., Gentili, L., Perfumi, M., De Caro, G. and Maggi, C.A., The tachykinin NH_2-senktide, a selective neurokinin B receptor agonist, is a very potent inhibitor of salt appetite in the rat, 341
McCart, R. and Straznicky, C., Lack of axon regeneration of isthmic neurons in juvenile *Xenopus*, 143
McIntyre, I., see Bhattacharya, S.K., 218
McRae-Degueurce, A., Hjorth, S., Dillon, D.L., Mason, D.W. and Tice, T.R., Implantable microencapsulated dopamine (DA): a new approach for slow-release DA delivery into brain tissue, 303
Meli, A., see Renzi, D., 254
Meli, A., see Santicioli, P., 197
Minami, M., see Kuraishi, Y., 325
Misbahuddin, M., Oka, M., Nakanishi, A. and Morita, K., Stimulatory effect of vasoactive intestinal polypeptide on catecholamine secretion from isolated guinea pig adrenal chromaffin cells, 202
Miyadai, T., see Hirayama, M., 34
Mizuno, Y., see Saito, T., 321
Moguilewsky, M., see Sarrieau, A., 14
Morgan, P.F., see Daval, J.L., 82
Morita, K., see Misbahuddin, M., 202
Murzi, E., see Baptista, T., 161

Nagatsu, T., see Saito, T., 321
Nagel, R.G., Busch, K.T. and Lee, H.-Y., A morphometric study of cytoskeletal reorganization in rat sciatic nerve axons following β,β'-iminodipropionitrile (IDPN) treatment, 241
Nakai, Y., see Iwase, M., 30
Nakajima, T., see Daval, J.L., 82
Nakanishi, A., see Misbahuddin, M., 202
Nanayama, T., see Kuraishi, Y., 325
Neff, N.H., see Weihmuller, F.B., 207
Neuhuber, W.L., see Doerffler-Melly, J., 121
Neuman, R.S., Ben-Ari, Y., Gho, M. and Cherubini, E., Blockade of excitatory synaptic transmission by 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) in the hippocampus in vitro, 64
Nicole, A., see Delacourte, A., 247
Nixon, R.A., see Shea, T.B., 291
Notter, M.F.D., see Del Cerro, M., 21

Ohno, H., see Kuraishi, Y., 325
Oka, M., see Misbahuddin, M., 202
Olsson, T., see Maehlen, J., 8
Ong, J., see Kerr, D.B., 92
Orsingher, O.A., see Ramírez, O.A., 275
Oxenkrug, G., see Bhattacharya, S.K., 218

Parada, M.A., see Baptista, T., 161
Pasquier, C., see Jacquin, T., 285
Pavlides, C., Greenstein, Y.J. and Winson, J., Comparative extracellular current flow at dendrites and soma of dentate granule cells during long-term potentiation, 177
Pellitteri, R., see Leanza, G., 132
Perfumi, M., see Massi, M., 341
Philibert, D., see Sarrieau, A., 14
Pifl, Ch., Schningnitz, G. and Hornykiewicz, O., The neurotoxin MPTP does not reproduce in the rhesus monkey the interregional pattern of striatal dopamine loss typical of human idiopathic Parkinson's disease, 228
Polidori, C., see Massi, M., 341
Pradelles, P., see Renzi, D., 254
Prager, R.H., see Curtis, D.R., 97
Prager, R.H., see Kerr, D.B., 92

Quadri, L., see Hösli, L., 165

Racké, K., see Wammack, R., 114
Ramírez, O.A., Orsingher, O.A. and Carrer, H.F., Differential threshold for long-term potentiation in the hippocampus of rats with inborn high or low learning capacity, 275
Regenold, J.T., Haas, H.L. and Illes, P., Effects of purinoceptor agonists on electrophysiological properties of rat mesencephalic trigeminal neurones in vitro, 347
Reiter, R.J., see Gonzalez-Brito, A., 330

Renzi, D., Santicioli, P., Maggi, C.A., Surrenti, C., Pradelles, P. and Meli, A., Capsaicin-induced release of substance P-like immunoreactivity from the guinea pig stomach in vitro and in vivo, 254
Roncada, P., see Gandolfi, O., 192
Rostène, W., see Sarrieau, A., 14
Russo, A., see Leanza, G., 132

Saito, H., see Tanaka, J., 187
Saito, T., Mizuno, Y., Nagatsu, T. and Yoshida, M., Effect of long-term administration of 1,2,3,4-tetrahydroisoquinoline (TIQ) on striatal dopamine and 3,4-dihydroxypenylacetic acid (DOPAC) content in mice, 321
Sandler, M., see Bhattacharya, S.K., 218
Santicioli, P., Maggi, C.A., Geppetti, P., Del Bianco, E., Theodorsson, E. and Meli, A., Release of calcitonin gene-related peptide-like immunoreactivity (CGRP-LI) from organs of the genitourinary tract in rats, 197
Santicioli, P., see Renzi, D., 254
Sarrieau, A., Dussaillant, M., Moguilewsky, M., Coutable, D., Philibert, D. and Rostène, W., Autoradiographic localization of glucocorticosteroid binding sites in rat brain after in vivo injection of [^{3}H]RU 28362, 14
Sato, K., see Hirayama, M., 34
Satoh, M., see Kuraishi, Y., 325
Satoh, M., see Ge, M., 76
Schneider, J.A., see Tsai, C., 298
Schningnitz, G., see Pifl, Ch., 228
Schröder, H.D., see Maehlen, J., 8
Seaton, P., see Doherty, P., 222
Seto, K., see Tanaka, J., 187
Shea, T.B., Majocha, R.E., Marotta, C.A. and Nixon, R.A., Soluble, phosphorylated forms of the high molecular weight neurofilament protein in perikarya of cultured neuronal cells, 291
Shioda, S., see Iwase, M., 30
Sinet, P.M., see Delacourte, A., 247
Smith, M.E., see Hughes, S., 1
Staiger, V., see Bonhoeffer, T., 259
Stanzani, S., see Leanza, G., 132
Straznicky, C., see McCart, R., 143
Stringer, J.L. and Lothman, E.W., NMDA receptor dependent paroxysmal discharges in the dentate gyrus, 69
Such, G., see Erdélyi, L., 46
Surrenti, C., see Renzi, D., 254

Tamura, A., see Kunugi, Y., 315
Tanabe, N. and Kijima, H., Transmitter release at frog end-plate loaded with a Ca^{2+}-chelator, BAPTA: hypertonicity and erythrosin B augment the release independently of internal Ca^{2+}, 52
Tanaka, J., Saito, H. and Seto, K., Involvement of the septum in the regulation of paraventricular vasopressin neurons by the subfornical organ in the rat, 187
Tannenbaum, M.G., see Gonzalez-Brito, A., 330
Theodorsson, E., see Santicioli, P., 197
Tice, T.R., see McRae-Degueurce, A., 303
Tsai, C., Schneider, J.A. and Lehmann, J., *Trans*-2-carboxy-3-pyrrolidineacetic acide (CPAA), a novel agonist at NMDA-type receptors, 298

Ueda, H., see Ge, M., 76

Vlahakos, D. and Gavras, H., Central sympathetic stimulation produced by saline application in the nucleus tractus solitarii area of conscious rats, 335

Walsh, F.S., see Doherty, P., 222
Wammack, R. and Racké, K., Endogenous opioid inhibition of the release of oxytocin from the isolated rat neurohypophysis during high-frequency stimulation of the pituitary stalk, 114
Weight, F.F., see Aguayo, L.G., 155
Weihmuller, F.B., Hadjiconstantinou, M., Bruno, J.P. and Neff, N.H., Administration of GM_1 ganglioside eliminates neuroleptic-induced sensorimotor deficits in MPTP-treated mice, 207
White, D.M. and Zimmerman, M., The bradykinin-induced release of substance P from nerve fibre endings in the rat saphenous nerve neuroma is not related to electrophysiological excitation, 108
White, S.R., see Jackson, D.A., 171
Wiegand, S.J., see Del Cerro, M., 21
Winson, J., see Pavlides, C., 177

Yang, J., see Zorumski, C.F., 86
Yinon, U., Chen, M., Zamir, S. and Gelerstein, S., Corpus callosum transection reduced binocularity of cells in the visual cortex of adult cats, 280
Yokochi, T., see Hirayama, M., 34
Yoshida, M., see Saito, T., 321

Zamir, S., see Yinon, U., 280
Zeevalk, G.D. and Hydman, A.G., Transferrin concentration and location during formation of chick retina: development correlates, 149
Zimmerman, M., see White, D.M., 108
Zorumski, C.F. and Yang, J., Non-competitive inhibition of GABA currents by phenothiazines in cultured chick spinal cord and rat hippocampal neurons, 86